Ketoconazole & Zinc pyrithione

DANDROP SHAMPOO



1. NAME OF THE MEDICINAL PRODUCT Ketoconazole & Zinc pyrithione

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

3. PHARMACEUTICAL FORM

Shampoo

4. CLINICAL PARTICULARS

4.1 Therapeutic Indications

Topical treatment of Seborrhoeic Dermatitis including dandruff resulting from proven fungal infection including Pityriasis versicolor.

4.2 Method of Administration

SHAKE WELL BEFORE USE

Moisten the hair and scalp thoroughly with water. Apply sufficient lotion to produce enough lather to wash scalp and hair and gently massage it over the entire scalp area for about 1 minute. Rinse hair thoroughly with water.

Repeat the procedure leaving the lotion on scalp for 3minutes. After the second thorough rinse, dry hair with towel or warm air flow. Use twice a week for 4 weeks with at least 3 days between each application and then intermittently as needed to maintain control or use as directed by physician.

The infected parts of the skin should be washed with DANDROP Shampoo 2%, allowing the product to soak in for 3 to 5 minutes before rinsing. The skin itself should be thoroughly washed, not just the hair.

4.3 Contraindications

DANDROP Shampoo 2% is contraindicated in individuals with unknown hypersensitivity to any of its ingredients.

4.4 Special Warnings and Special Precautions for Use

FOR EXTERNAL USE ONLY

In patients who have been on prolonged treatment with topical corticosteroids, it is recommended that the steroid therapy be gradually withdrawn over the period of 2 to 3 weeks, while using DANDROP Shampoo 2%, to prevent any potential rebound effect.

Avoid contact with the eyes. If the shampoo gets into the eyes, wash the eyes with copious amount of water.

Interactions with other Medicaments and other forms of Interaction: None known.

4.5 Fertility, Pregnancy and Lactation

There are no adequate and well-controlled studies in pregnant or lactating women. Plasma concentrations of ketoconazole were not detectable after topical administration of Ketoconazole Shampoo 2% to the scalp of non- pregnant humans.

There are no known risks associated with the use of DANDROP Shampoo 2% in pregnancy or lactation.

Adults, Elderly and Children:

'Use the shampoo twice weekly for 2-4 weeks. Wash affected areas of skin (not just the hair), using a palm full of shampoo and leave for 3-5 minutes before rinsing.

4.8 Undesirable Effects

Application site erythema, irritation, hypersensitivity, pruritus & pustules may occur

Others: Acne, Alopecia, Contact Dermatitis, Dry skin & hair texture, Abnormal skin rash and Skin burning sensation, Skin exfoliation, Unknown Angioedema, Urticaria, Hair color changes

4.9 Overdose

In the event of accidental ingestion, supportive and symptomatic measures should be carried out. To avoid aspiration, emesis or gastric lavage should be performed.

5. PHARMACOLOGICAL PROPERTIES

5.1 **Pharmacodynamic Properties**

Ketoconazole, a synthetic imidazole dioxolane derivative, has a potent antifungal activity against

dermatophytes, such as Trichophyton spp., Epidermophyton spp., Microsporum spp., and yeasts, such as

Candida spp. and Malassezia spp. (Pityrosporum spp.).

Zinc pyrithione is best known for its use in treating dandruff and seborrhoeic dermatitis. It has

fungistatic (that is, it inhibits the division of fungal cells) and bacteriostatic (inhibits bacterial cell

division) properties and is used in the treatment of seborrhoeic dermatitis. It is effective against

many pathogens from the Streptococcus and Staphylococcus genera.

DANDROP Shampoo 2% rapidly relieves scaling and pruritus, which are usually associated with

seborrhoeic dermatitis, pityriasis capitis (dandruff) and pityriasis versicolor.

6. PHARMACEUTICAL PARTICULARS

6.1 Shelf-life: 24 months

6.2 Special Precautions for Storage: Store below 25°C. Store in a cool and dry place protect from heat and

Light

6.3 Nature and contents of container: HDPE bottle

Marketed By:

Alkem Labs Limited

Alkem House;

SenapatiBapat Marg,

Lower Parel; Mumbai,

Maharashtra: 400 013.